



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

SPECIAL ARTICLES

ORGANIZATION OF HETEROTYPIC CHROMOSOMES

IN a recent article H. C. Sands¹ gives a preliminary account of his interpretation of chromosome structure in *Tradescantia*. This account seems to differ much in fundamentals from the results of a study conducted by the writer on various species of the South African Liliaceous genus *Gasteria*. To consider only the most interesting feature, the heterotypic chromosomes, we find first that at metaphase there are three pairs of small chromosomes in the center of the plate, with four pairs of much larger ones arranged in radiating fashion around them. In favorable material it is easy to see that the metaphase chromosomes are organized internally into chromomeres, though these are somewhat irregularly arranged. Even before they separate the larger chromosomes each show a cleft at the outer end opposite the fiber attachment, and before anaphase is far advanced they are nearly completely divided longitudinally into two halves. Usually this is entirely completed before telophase sets in, and the halves are widely separated. The smaller chromosomes are slow to split, but finally do so. Consequently there is at telophase a diploid number of separate chromosome-halves and the homotypic division has been prepared for.

As the larger chromosomes separate in anaphase it is clear that in each half-chromosome there is a double row of rounded chromomeres. These are few and apparently quite definite in number, well separated from each other and stain sharply in contrast to the almost colorless matrix. As the membrane first appears around the group of telophase chromosomes (which are well separated in this genus and easily studied) it becomes evident that there are no longer two rows of chromomeres, but that four are now present. When the chromosomes are obliquely placed these can clearly be distinguished, especially in the later stages when the chromosomes begin to broaden at the ends in preparation for a more even distribution of the chromatin in the interkinetic nucleus. The smaller roundish chromosomes

show comparable changes, though less clearly because of their size. It would seem that this can only mean that both the gametophyte divisions in the pollen grain have been completely prepared for. The chromomeres which enter the construction of the sperm nuclei seem to be already formed and merely awaiting the mechanical distribution of these mitoses. Unfortunately it has not been possible to trace the history of the chromomeres as such through these two divisions.

By a method based on smear preparations it has been possible to largely confirm the results of sectioned material and to obtain fixations in which synizesis (synapsis) has been almost entirely eliminated. The best results on the anaphase and telophase stages were obtained when the dividing cells were pressed from the anther before fixing and embedding.

The writer would consider, then, that the material worked upon, *Gasteria*, tends to demonstrate a very great precision in the chromomere constitution of the meiotic chromosomes. It is hoped that it will be possible soon to make a full report giving the evidence for the views here expressed.

WM. RANDOLPH TAYLOR

UNIVERSITY OF PENNSYLVANIA

THE INTELLIGENCE OF INDIANS

THE accompanying tables show that if we rank mixed bloods of certain heredity with nomadic and sedentary full blood Indians according to intelligence as indicated by Scale A of National Intelligence Tests the sequence proves to be:

- I. Mixed Bloods.
- II. Mexicans.
- III. Plains and South Eastern Full Bloods.
- IV. Plateau Indians, Full Bloods.
- V. Navajos and Apaches, Full Bloods.

While the number of cases is small the indications of the measures are consistent with this inference. The results will shortly be reported more in detail by the writer who has been giving these tests in United States Indian Schools of the Southwest.

THE RELATIVE INTELLIGENCE OF INDIANS OF NOMADIC AND SEDENTARY TRIBES AND MIXED BLOOD INDIANS

The scores are of the National Intelligence Tests, Scale A, Form 1.

¹ Sands, H. C.: "Perigenesis," SCIENCE LVI, 517-518, 1922.

	No. Cases	Score Median	Per cent. attaining Median of Pl. & S.E.
<i>12 and 13 years:</i>			
Mixed Bloods.....	15	103	80 per cent.
Mexicans	145	85	60 per cent.
Plains & S. E.....	8	76	—
Pueblo	46	64	40 per cent.
Navajo & Apache.....	12	52	10 per cent.
<i>14 and 15 years:</i>			
Mixed Bloods.....	39	110	80 per cent.
Mexicans	132	92	70 per cent.
Plains & S. E.....	55	85	—
Pueblo	82	80	44 per cent.
Navajo & Apache.....	19	60	20 per cent.
<i>16 and 17 years:</i>			
Mixed Bloods.....	41	104	71 per cent.
Mexicans	28	91	54 per cent.
Plains & S. E.....	60	90	—
Pueblo	95	78	34 per cent.
Navajo & Apache.....	30	77	23 per cent.
<i>18 and 19 years:</i>			
Mixed Bloods.....	31	114	60 per cent.
Mexicans	2	—	—
Plains & S. E.....	53	88	—
Pueblo	26	71	30 per cent.
Navajo & Apache.....	24	77	40 per cent.

T. R. GARTH

UNIVERSITY OF TEXAS
JULY 8, 1922.THE OPTICAL SOCIETY OF
AMERICAABRIDGED MINUTES OF THE SEVENTH
ANNUAL MEETING¹

THE seventh annual meeting of the Optical Society of America was held at the National Bureau of Standards, Washington, on October 25, 26, 27 and 28, President Troland presiding.

One hundred and thirteen persons registered as in attendance at the meeting. Eighty-five of these were from outside Washington. The actual number attending at one time or another was probably about 150. The number in attendance at any one time at the sessions for the reading of papers ranged from about thirty to over one hundred.

¹ The complete minutes including abstracts of papers and descriptions of exhibits will appear in the January number of the *Journal of the Optical Society of America*.

An exhibit of optical instruments was held in connection with this meeting; and visitors were also given an opportunity to inspect the optical equipment of the Bureau of Standards. The condensed program follows:

OCTOBER 25

Bureau of Standards laboratory exhibits open to informal visits.

Business meeting:

Reports of officers and general committees.

Reports of committees on nomenclature and standards.

OCTOBER 26

Glass Plant open to informal visits. Pot of optical glass being stirred.

General session:

Address of welcome: Dr. S. W. Stratton, director, Bureau of Standards.

Response: Dr. Leonard T. Troland, president, Optical Society.

Papers on miscellaneous optics.

Session on solar, stellar and planetary radiation: Invited papers by Dr. C. G. Abbot and Dr. W. W. Coblentz.

Glass Plant open to informal visits.

Moulding, annealing and inspection of optical glass.

Illustrated address (by invitation): Professor W. J. Humphreys, U. S. Weather Bureau, on "The optics of the atmosphere."

Glass Plant open to visitors:

Pot of optical glass removed from furnace.

OCTOBER 27

First session on physiologic optics.

Second session on physiologic optics.

Special session on physiologic optics for papers by invitation of the committee.

OCTOBER 28

Session on photometry, colorimetry and optical pyrometry.

The exhibit of optical instruments, October 26, 27 and 28.

A synopsis of the proceedings, papers and special features of the meeting is given below:

1. *Business*

The report of the secretary and membership committee was read in part.

The president communicated an informal report from the treasurer, the formal report to be submitted at the end of the fiscal year.

The following reports of general committees were received informally:

Physiologic optics: F. K. Richtmyer, chairman.